



Phytodiversity of Barail Wildlife Sanctuary, Assam, India: field-based observations—I. Trees and lianas

Arpita Bora, Debjyoti Bhattacharyya

Plant Taxonomy and Biosystematics Laboratory, Department of Life Science and Bioinformatics, Assam University, Silchar 788 011, Assam, India.
Corresponding author: Debjyoti Bhattacharyya, dbhattacharyya_au@yahoo.in

Abstract

Trees and lianas were inventorized in Barail Wildlife Sanctuary, Assam, India, which revealed the occurrence of 112 and 23 species of trees and lianas respectively. Two separate lists, one for trees and other for lianas, of all the enumerated taxa are presented here with their vernacular names, phenology, places of occurrence and collection number(s). Threat status of each taxon according to IUCN Red List categories is also provided. Anthropogenic activities and other natural calamities are causing serious threats particularly to the tree diversity of the sanctuary. In addition, destruction of one tree may eradicate many lianas from a particular area. Thus, many species of trees and lianas are under alarming threat in the sanctuary. Conservation through *in situ* and *ex situ* modi operandi is recommended.

Key words

Arboreal angiosperms; Barak Valley; biodiversity; Cachar; species inventory; woody climbers.

Academic editor: Amit Kumar | Received 9 May 2017 | Accepted 22 September 2017 | Published 15 December 2017

Citation: Bora A, Bhattacharyya D (2017). Phytodiversity of Barail Wildlife Sanctuary, Assam, India: field-based observations—I. Trees and lianas. Check List 13 (6): 1037–1053. <https://doi.org/10.15560/13.6.1037>

Introduction

Inventorization of plants of a particular area comprises basic biodiversity studies and assessment of their conservation status. Among various ecoclimatic zones, tropical evergreen forests constitute one of the most biodiversity rich habitats on the globe (Khandekar and Srivastava 2014). An exhaustive inventory of tropical plant species is, thus, one of the primary objectives for the plant taxonomists. Within India, the tropical monsoon climate that prevails in the Eastern Himalayas, Northeastern India, and Western Ghats allows for a rich and diverse flora. Though steps have been taken towards cataloguing the biodiversity of protected areas in India, knowledge of the diverse flora of many protected areas, particularly in northeastern India, is still lacking. There was no comprehensive study on the diversity of angiosperms of Barail

Wildlife Sanctuary (BWS), Assam, India, and hence, we initiated this study to investigate the same of BWS for the first time. Because the angiosperm flora of BWS is large, we have segregated the study into 2 parts. A second part, which will include herbs, shrubs, and non-woody climbers, is planned for a later publication.

Tropical vegetation, mostly dominated by angiosperm trees, is a large part of floral diversity. Trees are important in that they provide shade and shelter for herbaceous and epiphytic plants. They provide homes for wild animals, especially primates. They stabilise the soil, counteract rapid run-off of rainwater, and prevent desertification, and they have a role in climate control and in the maintenance of biodiversity and ecosystem balance (Bellefontaine et al. 2002). Trees are also economically important for yielding timber, fuel, and food.

Lianas, or woody climbers, are likewise important components of tropical forests and are important for the forest community and structure. They are generally long-stemmed, reaching up into the tree canopy to get access to light. Their structural morphology and mechanisms of twining add to the complexity of the vertical structures of forests (Chaudhuri and Naithani 1985). According to Grubb (1977), dense vegetation of lianas in a particular area portray the nature of tropical lowland and lower montane forest types of that region. Like trees, lianas also contribute substantially to the diversity of the forests and provide food and arboreal pathways for forest animals. They are widely used for medicine, food, house construction, and artisan work. They have an important role in forest regeneration (Bongers et al. 2002).

The first comprehensive floral account of Assam was the *Flora of Assam* (Kanjilal et al. 1934–1940). Subsequently, Das (1942) documented preliminary data on flora of Assam. Recently, Choudhury (2005) and Barooah and Ahmed (2014) provided checklists of the vascular plants, particularly angiosperms and gymnosperms, of the state. Additionally, smaller works on the flora in various areas of the state, particularly from the Brahmaputra Valley region, include: Rao and Rabha (1966), Rao and Verma (1970, 1972, 1973, 1976), Islam (1986, 1990, 1990a, 1991), Sarma (1990), Barua (1992), Choudhury et al. (1994), Agarwal and Borah (2001), Barooah and Mahanta (2006), Sarma et al. (2006).

By comparison, the Barak Valley (southern Assam), where Barail Wildlife Sanctuary is situated, has received very little attention by botanists. Dutt et al. (1974) prepared a preliminary checklist of the Polypetalae of North Cachar Hills and Barail range (particularly the Haflong hills). Other reports from Barak Valley were made by Sharma et al. (2002), Das and Dutta Choudhury (2003), Nath and Maiti (2003), Baruah et al. (2006), Das et al. (2006, 2007, 2008), Barbhuiya et al. (2009), Darlong and Bhattacharyya (2011), Rout et al. (2012), Darlong et al. (2013), Barbhuiya 2013a, Devi and Bhattacharyya (2013, 2013a, 2013b, 2014, 2016), Barbhuiya 2014, Barbhuiya et al. (2014), and Devi et al. (2014). Das et al. (2013) and Baruah and Dutta Choudhury (2015) listed the herbaceous plants of Barak Valley, focusing on their economic utility.

Among the angiosperms of the BWS, while describing a new Asian genus of ginger, *Larsenianthus*, Kress et al. (2010) reported 2 species of the genus from the sanctuary viz. *L. assamensis* S. Dey, Mood & S. Choudhury (described as new to science from BWS) and *L. careyanus* (Benth.) W.J. Kress & Mood. Barbhuiya et al. (2012) and Barbhuiya (2013) described 2 new species, *Ornithochilus cacharensis* Barbhuiya, B.K. Dutta & Schuit. (Orchidaceae) and *Toxicodendron bimannii* Barbhuiya (Anacardiaceae), respectively. Barbhuiya et al. (2012a) rediscovered *Ardisia keenanii* C.B. Clarke (Myrsinaceae) after a gap of 138 years from the sanctuary. First collection of this species from the study area was made by R.L. Keenan in 1873. Later, C.B. Clarke (1882) described it in *Flora of British India* based on his own collection from

Manipur in 1868 and Keenan's collection from Cachar in 1873. Barbhuiya et al. (2013) documented 96 species of grasses from Barak Valley, including 16 species from the BWS.

As can be seen from the literature, it is evident that baseline data on angiosperm diversity of the BWS is mostly scattered and inadequate. Hence, we initiated this study to provide a comprehensive list of angiosperms in the sanctuary. In this first part, we treat the trees and lianas of the protected area.

Methods

Study site. Barail Wildlife Sanctuary (BWS) is situated in the northern part of Cachar district of Assam and lies along the foothills of the North Cachar and Barail hills. It covers an area of just over 326 km². The BWS is under the administrative control of the Southern Assam Forest Circle, Silchar, and consists of Barail Reserve Forest, which is part of the Cachar Forest Division (East Block) and North Cachar Reserve Forest, part of the Karimgunj Forest Division (West Block) (Fig. 1).

The climate is tropical and humid; the average annual rainfall is 3383.5 mm/year and the average humidity is 72–83%. The minimum and maximum temperatures recorded during our study were 12.6 °C and 32.7 °C, respectively.

Major rivers draining the sanctuary are the Jatinga, Daloo, Kayong, Gumra, and Boleswar. The sanctuary is characterized by undulating hills having altitudinal range of less than 30 m to more than 1800 m in the Barail hills, where the towering peaks are Hamplopit (1,867 m), Kua-kaha (1,736 m), Maheo (1,739 m), and Sherpai (1,657 m) (Anonymous 2006). Geologically, the soils are sandy-stony to clayey (Anonymous 2006). The area is one of the richest treasure houses of flora as well as fauna due to its unique geographical position, diverse landscapes, wide range of physiographic conditions and high precipitation.

As per classification of Champion and Seth (1968), the sanctuary has 2 broad groups of forests, tropical wet evergreen and tropical semi-evergreen forests.

Field survey and data collection. We made extensive and intensive field observations and collections from 2013 to 2016 (Table 1), from various habitats and forest types (Figs 2, 3). Our surveys and collections were done with the permission from the Divisional Forest Officer, Cachar Division, Silchar (letter No.B/35/Misc.7344-45, dated Silchar, 21 November 2013). Although we surveyed in almost all seasons, most field surveys were done during the pre-monsoon and post-monsoon seasons. Specimens were collected mostly during their flowering and fruiting stages. Vernacular names, phenological data, and ecological notes were compiled from our field observations and a survey of the literature; these data were supplemented by collectors' notes on herbarium sheets. The IUCN Red List was used to categorize species (IUCN 2017) into different Red List categories.




Figure 1. Map of the Barail Wildlife Sanctuary in Assam, India. See Table 1 for numbered localities.

Table 1. Localities, with dates of collections (surveys were 4–5 days at each location), elevation, and geographic coordinates. Locality number corresponds to numbers in Figure 1.

| No. | Location | Date | Elevation (m) | Latitude (N) | Longitude (E) |
|--------|--|---------------------------------------|---------------|---------------|-----------------|
| 1 | Bandarkhal and vicinity | 24-IX-2013 | 94–102 | 25°16′ | 092°51′ |
| 2 | Craig Park Tea Estate, Kalain | 26-XI-2014 | 19–20 | 25°00′ | 092°52′ |
| 3 | Damcherra and vicinity | 11-III-2013, 12-XII-2013, 12-XII-2014 | 23–67 | 25°01′ | 092°49′ |
| 4 | Ditekcherra | 12-XII-2013 | 92–105 | 25°00′ | 092°51′ |
| 5 | Durbintila (on way to peak from foothills) | 12-XII-2013, 08-III-2014, 26-II-2016 | 105–1050 | 24°95′ | 092°51′ |
| 6 | Indranagar | 18-V-2013 | 22–65 | 24°55′ | 092°52′ |
| 7 | Jatingacherra and vicinity | 28-II-2015 | 965–970 | 25°50′ | 092°49′ |
| 8 | Kalain Tea Estate, hill no. 15 | 26-XI-2014 | 78–85 | 24°55′ | 092°47′ |
| 9 | Kumba | 02-IV-15 | 69–75 | 24°94′ | 092°52′ |
| 10 | Lakhicherra | 24-IV-2014 | 105–110 | 24°58′ | 092°50′ |
| 11 | Madhura Khuwari | 05-IV-2013, 07-IV-2013 | 25–35 | 26°11′ | 091°83′ |
| 12 | Malidahar | 03-III-2013, 12-XII-2015 | 15–25 | 25°03′ | 092°38′ |
| 13, 14 | Malidahar Mikir punji and vicinity | 12-XII-2015 | 19–30 | 25°01′–25°08′ | 092°48′–092°59′ |
| 15 | Maruacherra | 08-III-2014 | 105 | 24°99′ | 092°50′ |
| 16 | SCF Nala, on way to Damcherra | 25-IV-2015 | 103–135 | 24°59′ | 092°45′ |




Figure 2. Forest habitats at various locations of Barail Wildlife Sanctuary. **A.** SFC Nala. **B.** Durbintila. **C.** Indranagar. **D.** Malidahar. **E.** Confluence of Kayang and Jatinga River. **F.** Lakhicherra.




Figure 3. Forest environments of the study area. **A–C.** Dense forests in core areas. **D–F.** Liana dominated forests.

Processing of specimens. The drying, treatment with chemicals, and mounting of our specimens were done following the routine herbarium practices (Jain and Rao 1977, Singh and Subramaniam 2008). Voucher specimens were deposited in the Plant Taxonomy and Biosystematics laboratory, Department of Life Science and Bioinformatics, Assam University, Silchar (here abbreviated as AULS; Tables 2, 3).

Identification of species. We studied the external morphological characteristics, including those of dissected, mainly floral parts, of all collected specimens using an Olympus SZ61 stereomicroscope. Characteristics of each species and infraspecific taxon were noted. Preliminary identifications were made using keys in authoritative literature (Brandis 1906, Kanjilal et al. 1934–1940, Chaudhuri and Naithani 1985). We confirmed the identifications of our specimens with authentic herbarium specimens housed in Indian herbaria (ASSAM, BSD, CAL, DD, GUBH, and LWG; acronyms according to Thiers 2011) and with images of type specimens available in online databases. The *International Plant Name Index* (IPNI 2012), *The Plant List* (2013) and *Tropicos* (2017) were consulted for current nomenclature of all taxa. The Angiosperm Phylogeny Group III Classification (APG III 2009) was followed for the classification of families. Families and species in our list are arranged in alphabetical order.

Results

Floristic diversity. Our study revealed 112 species of trees belonging to 86 genera and 43 families (Table 2), and 23 species of lianas in 17 genera and 13 families (Table 3).




Figure 4. Dominant tree families with number of species and genera.

Of the species of trees, among the dicotyledons, the most dominant families were Fabaceae and Moraceae with 10 species each, followed by Euphorbiaceae (8), Meliaceae (7), Lauraceae (6), and Rubiaceae and Rutaceae (4 each) (Fig. 4). Seven families—Anacardiaceae, Apocynaceae, Bignoniaceae, Combretaceae, Elaeocarpaceae, Malvaceae, and Myrtaceae—each had 3 species. Actinidiaceae, Boraginaceae, Dilleniaceae, Lamiaceae, Primulaceae, Rhamnaceae, and Verbenaceae had 2 spp. each. The remaining 20 families were represented by a single species in the BWS. *Ficus* (6 spp.), followed by *Artocarpus* (4 spp.), *Bauhinia*, *Litsea*, and *Terminalia* (3 spp. each) were the dominant genera in terms of species numbers. Among the monocotyledons, only Poaceae (5 spp.) and Arecaceae (3 spp.) were recorded. Bamboos (Poaceae) and Areca nut (Arecaceae) were 2 dominant floral components in parts of the BWS.

Of the lianas, among the dicotyledons, the most dominant families were Fabaceae (8 spp.), followed by

Table 2. List of trees with their vernacular names, phenology, collection number, occurrence and threat status. Abbreviations, languages: A = Assamese, B = Bengali, Di = Dimasa, E = English, H = Hindi, K = Kuki, Sylh. = Sylheti; IUCN status: DD = Data Deficient, LC = Least Concern, NE = Not Evaluated and V = Vulnerable. Occurrence: see Table 1 for a key to locations.

| Scientific name | Vernacular name | Phenology | Voucher specimens | Occurrence | Threat status |
|--|------------------------------------|------------|--------------------------|-------------|---------------|
| Dicotyledons | | | | | |
| Actinidiaceae Engl. & Gilg | | | | | |
| <i>Saurauia cerea</i> Griff. ex Dyer | Not known | July–Nov. | AB & DB 11479; AULS 2554 | 10 | NE |
| <i>Saurauia roxburghii</i> Wall. | Arbeng-thing (B) | March–Aug. | AB & DB 11606; AULS 2560 | 11 | NE |
| Anacardiaceae R.Br. | | | | | |
| <i>Lannea coromandelica</i> (Houtt.) Merr. | Kuhimola (A) | Feb.–June | AB & DB 11518; AULS 2551 | 16 | NE |
| <i>Mangifera indica</i> L. | Aam (A and B) | Year round | AB & DB 11554; AULS 2557 | 1, way to 7 | DD |
| <i>Spondias pinnata</i> (L.f.) Kurz | Amora (A) | March–Dec. | AB & DB 11555; AULS 2563 | way to 1 | NE |
| Annonaceae Juss. | | | | | |
| <i>Annona squamosa</i> L. | Atlash kothal (A), Sugar apple (E) | June–Sept. | AB & DB 11556; AULS 2568 | 1 | NE |
| Apocynaceae Juss. | | | | | |
| <i>Alstonia scholaris</i> (L.) R.Br. | Sotiona (A), Chhatim (B) | June–Dec. | AB & DB 11515; AULS 2556 | 1, way to 7 | LC |
| <i>Holarrhena pubescens</i> Wall. ex G.Don | Dhulkari (A), Kurchi (B) | Apr.–Dec. | AB & DB 11734; AULS 2571 | 4 | LC |
| <i>Plumeria rubra</i> L. | Goalanchi (A) | June–Dec. | AB & DB 11558; AULS 2562 | 1 | NE |
| Araliaceae Juss. | | | | | |
| <i>Trevesia palmata</i> (Roxb. ex Lindl.) Vis. | Snowflake Tree (E) | Oct.–July | AB & DB 11431; AULS 2580 | 5 | NE |
| Bignoniaceae Juss. | | | | | |
| <i>Oroxylum indicum</i> (L.) Kurz | Thukuna gach (B) | Jan.–March | AB & DB 11564; AULS 2585 | 5 | NE |

Table 2. Continued.

| Scientific name | Vernacular name | Phenology | Voucher specimens | Occurrence | Threat status |
|---|--|-------------|---------------------------------|---------------------------------|---------------|
| <i>Pajanelia longifolia</i> Schum | Wild Trumpet tree (E), Hona, Kawarinoa (Sylh.) | Dec.–May | AB & DB 11565; AULS 2588 | 5 | NE |
| <i>Stereospermum chelonoides</i> DC. | Parhori, Paroli (A), Fragrant Padri tree (E) | Apr.–Jan. | AB & DB 11761; AULS 2597 | 6 | NE |
| Boraginaceae Juss. | | | | | |
| <i>Cordia myxa</i> L. | Bowal (A), Indian cherry (E) | May–Dec. | AB & DB 11777; AULS 2582 | 16 | NE |
| <i>Ehretia acuminata</i> R.Br. | Bowal gach, gual, pojhar (A), Koda Tree (E) | March–Jan. | AB & DB 11524; AULS 2594 | 3 | NE |
| Burseraceae Kunth | | | | | |
| <i>Garuga pinnata</i> Roxb. | Bombuk, Engla (B) | March–Oct. | AB & DB 11569; AULS 2600 | 5 | NE |
| Caprifoliaceae Juss. | | | | | |
| <i>Viburnum cylindricum</i> Buch.-Ham. ex D.Don | Not known | June–Oct. | AB & DB 11413; AULS 2617 | 3 | NE |
| Capparaceae Juss. | | | | | |
| <i>Crateva religiosa</i> G.Forst. | Barun Gach (A) | March–Sept. | AB & DB 11771; AULS 2613 | 11 | NE |
| Caricaceae Dumort. | | | | | |
| <i>Carica papaya</i> L. | Omita (A), Papaya (E) | March–Dec. | AB & DB 11578; AULS 2568 | 1 | NE |
| Clusiaceae Lindl. | | | | | |
| <i>Mesua ferrea</i> L. | Nahor (A), Nageswar (H), Laternga (Jaintia) | March–Oct. | AB & DB 11755; AULS 2552 | 6 | NE |
| Combretaceae R.Br. | | | | | |
| <i>Terminalia bellerica</i> (Gaertn.) Roxb. | Bhumora (A), Bohera (B) | March–July | AB & DB 11756; AULS 2573 | 6 | NE |
| <i>Terminalia chebula</i> Retz. | Silikha (A), Haritoki (B), Sohartaki (Jaintia) | May–Dec. | AB & DB 11580; AULS 2623 | 1 | NE |
| <i>Terminalia myriocarpa</i> Van Heurck & Müll.Arg. | Hollokh (A), Jhalna (B) | Aug.–Jan. | AB & DB 11757; AULS 2601 | 6, 11 | NE |
| Datiscaceae Dumort. | | | | | |
| <i>Tetrameles nudiflora</i> R.Br. | Bhelu (A), Chundul (B) | March–May | AB & DB 11779; AULS 2610 | 16 | LC |
| Dilleniaceae Salisb. | | | | | |
| <i>Dillenia indica</i> L. | Ou tenga (A), Chalta (B) | July–Jan. | AB 11586; AULS 2628 | Kalaincherra stream (through 8) | NE |
| <i>Dillenia pentagyna</i> Roxb. | Akshi, Okshi (A) | Apr.–Sept. | AB & DB 11763; AULS 2619 | 11 | NE |
| Dipterocarpaceae Blume | | | | | |
| <i>Shorea robusta</i> C.F.Gaertn. | Sal (A) | Jan.–July | AB 11746; AULS 2605 | 12 | LC |
| Elaeocarpaceae Juss. ex DC. | | | | | |
| <i>Sloanea sterculiacea</i> Rehder & E.H.Wilson var. <i>assamica</i> (Benth.) Coode | Joba-hingori, Bandor-kakoi (A) | Oct.–Apr. | AB & DB 11768; AULS 2644 | 11 | NE |
| <i>Elaeocarpus floribundus</i> Blume | Jalphai (A), Belphai (B) | May–Dec. | AB 11748; AULS 2651 | Kalaincherra stream (through 8) | NE |
| <i>Elaeocarpus lanceifolius</i> Roxb. | Not known | June–Sept. | AB 11504; AULS 2657 | Kalaincherra stream (through 8) | NE |
| Euphorbiaceae Juss. | | | | | |
| <i>Antidesma ghesaembilla</i> Gaertn. | Heloch, Mikhan-tenga (A), Black Currant Tree (E) | March–Jan. | AB & DB 11303; AULS 2564 | 3 | NE |
| <i>Aporosa octandra</i> (Buch.-Ham. ex D.Don) A.R.Vickery | Bara-heloch (A), Kokra (B) | Jan.–Dec. | AB & DB 11607; AULS 2553 | 11 | NE |
| <i>Bischofia javanica</i> Blume | Urim (A) | Apr.–Oct. | AB & DB 11461; AULS 2592 | 10 | NE |
| <i>Glochidion khasicum</i> Hook.f. | Not known | June–Nov. | AB & DB 11434; AULS 2599 | 5 | NE |
| <i>Glochidion lanceolarium</i> (Roxb.) Voigt | Not known | Apr.–Feb. | AB & DB 11480, 11497; AULS 2583 | 10 | NE |
| <i>Macaranga denticulata</i> (Blume) Mull.Arg. | Moralia (A), Jagura (B) | Apr.–Oct. | AB 11311, 11450; AULS 2575 | 5, 12 | NE |
| <i>Mallotus philippensis</i> (Lam.) Mull.Arg. | Monkey-face Tree (E), Kamala (H) | March–Aug. | AB & DB 11425; AULS 2602 | 5 | NE |
| <i>Phyllanthus emblica</i> L. | Amlakhi (A) | Apr.–Sept. | AB & DB 11749; AULS 2625 | 6 | NE |
| Fabaceae Lindl. | | | | | |
| <i>Bauhinia acuminata</i> L. | Kanchan (B) | Apr.–Dec. | AB & DB 11437; AULS 2674 | 5 | NE |
| <i>Bauhinia purpurea</i> L. | Kanchan (B) | Sept.–March | AB 11388, 11688; AULS 2684 | 2, 3 | LC |
| <i>Bauhinia variegata</i> L. | Kotora (A), Raktakanchan (B) | Dec.–Sept. | AB & DB 11740; AULS 2689 | on way to 1 | LC |
| <i>Cassia fistula</i> L. | Shonalu (B) | Apr.–Jan. | AB & DB 11733; AULS 2640 | 11 | NE |

Table 2. Continued.

| Scientific name | Vernacular name | Phenology | Voucher specimens | Occurrence | Threat status |
|--|--|-------------|---------------------------------|--------------------------|---------------|
| <i>Cassia javanica</i> L. subsp. <i>nodosa</i> (Buch.-Ham. ex Roxb.) K.Larsen & S.S.Larsen | Apple blossom cassia (E) | Aug.–Jan. | AB & DB 11547; AULS 2648 | 5 | NE |
| <i>Dalbergia sericea</i> Spreng. | Not known | Apr.–Sept. | AB 11604, 11684; AULS 2660 | 11, 12 | NE |
| <i>Erythrina variegata</i> L. | Indian coral tree (E) | Feb.–Aug. | AB & DB11732; AULS 2678 | 6, 11 | LC |
| <i>Pongamia pinnata</i> (L.) Pierre | Karchaw (A) | May–Oct. | AB & DB 11522; AULS 2661 | 16 | LC |
| <i>Saraca asoca</i> (Roxb.) de Wilde | Ashok (B) | March–July | AB & DB 11474; AULS 2672 | 10 | V |
| <i>Tamarindus indica</i> L. | Teteli (A), Tetul (B) | May–Apr. | AB 11576; AULS 2607 | 2 | NE |
| Lamiaceae Martinov | | | | | |
| <i>Callicarpa arborea</i> Roxb. | Bonmola/Gunmola/Khoja (A), Bormala (B), Beautyberry Tree (E) | May–Dec. | AB & DB 11531; AULS 2630 | 16 | NE |
| <i>Premna bengalensis</i> Clarke | Gohora (A) | May–Nov. | AB & DB 11760; AULS 2615 | 6, 11 | NE |
| Lauraceae Juss | | | | | |
| <i>Alseodaphne owdeni</i> R.Parker | Tilsundi (Cach.) | May | AB & DB 11775; AULS 2558 | 16 | NE |
| <i>Cinnamomum glanduliferum</i> (Wall.) Meisn. | Gonhorai (A) | March–Sept. | AB & DB 11762; AULS 2567 | 11 | NE |
| <i>Cinnamomum tamala</i> (Buch.-Ham.) T.Nees & C.H.Eberm. | Tej-pat (A) | Apr.–Oct. | AB & DB 11730; AULS 2608 | 11 | NE |
| <i>Litsea cubeba</i> Pers. | May Chang / Mountain Pepper (E) | Feb.–Aug. | AB 11502; AULS 2621 | Kalaincherra (through 8) | NE |
| <i>Litsea glutinosa</i> (Lour.) C.B.Rob. | Indian laurel (E) | May–Oct. | AB & DB 11625; AULS 2654 | 11 | NE |
| <i>Litsea monopetala</i> (Roxb.) Pers. | Sowalu (A) | May–Nov. | AB & DB 11447; AULS 2637 | 5 | NE |
| Lecythidaceae Poiteau | | | | | |
| <i>Barringtonia acutangula</i> (L.) Gaertn. | Hijal (A) | March–Oct. | AB & DB 11729; AULS 2643 | 11 | NE |
| Lythraceae J.St.-Hil. | | | | | |
| <i>Duabanga grandiflora</i> Walp. | Ramdala (B) | Feb.–June | AB & DB 11416; AULS 2647 | 5 | NE |
| Malvaceae Juss. | | | | | |
| <i>Bombax ceiba</i> L. | Simolu (A) | Jan.–May | AB 11566; AULS 2659 | 12 | NE |
| <i>Bombax insigne</i> Wall. | Bon Semal (B), Showy silk cotton tree (E) | Nov.–March | AB 11567; AULS 2667 | 2 | NE |
| <i>Sterculia versicolor</i> Wall. | Durong (A and Bodo) | March–June | AB & DB 11514; AULS 2688 | 1 | NE |
| Meliaceae Juss. | | | | | |
| <i>Aglaia spectabilis</i> (Miq.) S. S. Jain & Bennet | Amari, Bhoto-mayna (A) | Sept.–Nov. | AB & DB 11778; AULS 2690 | 16 | LC |
| <i>Aphanamixis polystachya</i> (Wall.) R. Parker | Not known | May–Apr. | AB & DB 11727; AULS 2555 | 3 | LC |
| <i>Azadirachta indica</i> Juss. | Neem (A) | March–July | AB & DB 11726; AULS 2572 | way to 5 | NE |
| <i>Chukrasia tabularis</i> A.Juss. | Boga poma, Hatia (A), Chikrassi (B) | June–Oct. | AB & DB 11758; AULS 2596 | 6 | LC |
| <i>Dysoxylum gotadhora</i> (Buch.-Ham.) Mabb. | Bandardima (A), Bandarfela (B) | March–Nov. | AB & DB 11759; AULS 2631 | 6 | NE |
| <i>Melia azedarach</i> L. | Ghura Neem (A) | March–Dec. | AB & DB 11453; AULS 2653 | 5 | NE |
| <i>Toona ciliata</i> M. Roem. | Poma (A), Red cedar (E) | Jan.–Aug. | AB & DB11454; AULS 2680 | 5 | LC |
| Moraceae Gaudich. | | | | | |
| <i>Artocarpus heterophyllus</i> Lam. | Kathal (A) | March–Aug. | AB & DB 11570; AULS 2686 | 5 | NE |
| <i>Artocarpus chama</i> Buch.-Ham. | Cham (A) | March–Aug. | AB & DB 11764; AULS 2663 | 6 | NE |
| <i>Artocarpus integra</i> (Thunb.) Merr. | Kathal (A, B) | Feb.–Sept. | AB & DB 11765; AULS 2676 | 11 | NE |
| <i>Artocarpus lacucha</i> Buch.-Ham. | Bohot (A), Dewa (B), Monkey Jack (E) | Feb.–Aug. | AB & DB 11766; AULS 2665 | 6 | NE |
| <i>Ficus benghalensis</i> L. | Bat (B) | Apr.–Nov. | AB & DB 11570; AULS 2569 | 11 | NE |
| <i>Ficus benjamina</i> L. | Jari gach (A) | Oct.–Apr. | AB & DB 11571; AULS 2561 | 3 | NE |
| <i>Ficus hispida</i> L.f. | Not known | June–July | AB & DB 11517, 11618; AULS 2574 | 11, 16 | NE |
| <i>Ficus palmata</i> Forssk. | Wild fig/Punjab Fig (E), Anjiri (H) | June–Sept. | AB & DB 11423, 11482; AULS 2591 | 5, 10 | NE |
| <i>Ficus racemosa</i> L. | Dumur (B) | Feb.–Sept. | AB & DB 11767; AULS 2598 | 6 | NE |
| <i>Ficus religiosa</i> L. | Anhat gach (A) | Apr.–Sept. | AB & DB 11572; AULS 2627 | 1 | NE |
| Moringaceae Martinov | | | | | |
| <i>Moringa oleifera</i> Lam. | Sojina (A), Drum Stick tree (E) | Year round | AB & DB 11725; AULS 2670 | Balacherra near 16 | NE |
| Myrtaceae Juss. | | | | | |
| <i>Psidium guajava</i> L. | Madhuriam (A), Guava (E) | Apr.–Oct. | AB & DB 11723; AULS 2683 | 6 | NE |
| <i>Syzygium cumini</i> (L.) Skeels | Jamuk (A), Jamun (B), Blackpalm (E) | Feb.–Sept. | AB & DB 11722; AULS 2578 | roadside at 1 | NE |
| <i>Syzygium kurzii</i> (Duthie) N. P.Balacr. | Bogi Jamuk (A) | Apr.–Oct. | AB & DB 11415; AULS 2590 | Roadside at 3 | NE |

Table 2. Continued.

| Scientific name | Vernacular name | Phenology | Voucher specimens | Occurrence | Threat status |
|---|---|-------------|--|--------------------------------------|---------------|
| Oleaceae Hoffmanns. & Link | | | | | |
| <i>Nyctanthes arbortritis</i> L. | Sewali (A), Shephalika (B) | Aug.–Jan. | AB 11720; AULS 2612 | Kalaincherra near 8 | NE |
| Primulaceae Batsch ex Borkh. | | | | | |
| <i>Ardisia paniculata</i> Roxb. | Not known | Year round | AB 11683; AULS 2629 | 12 | NE |
| <i>Maesa ramentacea</i> Wall. | Not known | Jan.–Dec. | AB & DB 11376; AULS 2646 | 4 (on way to 1) | NE |
| Rhamnaceae Juss. | | | | | |
| <i>Zizyphus jujuba</i> Mill. | Bogori (A) | May–Oct. | AB & DB 11706; AULS 2675 | 1 | LC |
| <i>Zizyphus rugosa</i> Lamk. | Suran (H), Bon-boroi (Sylh.) | Nov.–Jan. | AB 11705; AULS 2668 | 8 | NE |
| Rubiaceae Juss. | | | | | |
| <i>Chassalia curviflora</i> (Wall.) Thwaites var. <i>longifolia</i> Hook.f. | Not Known | May–July | AB & DB 11477, 11491; AULS 2559 | 10 | NE |
| <i>Morinda angustifolia</i> Roxb. | Not known | Apr.–Oct. | AB & DB 1481, 11627; AULS 2565 | 10, 11 | NE |
| <i>Neolamarckia cadamba</i> (Roxb.) Bosser | Kadam (A) | June–Nov. | AB & DB 11704; AULS 2616 | 1 | NE |
| <i>Wendlandia budleioides</i> Wall. ex Wight & Arn. | Not known | Aug.–Dec. | AB & DB 11424; AULS 2624 | 5 | NE |
| Rutaceae Juss. | | | | | |
| <i>Aegle marmelos</i> (L.) Correa | Bel (A and B) | Apr.–Aug. | AB & DB 11702; AULS 2587 | 10 | NE |
| <i>Citrus maxima</i> (Burm.) Merr. | Robab tenga (A), Chinese grapefruit (E) | Apr.–Dec. | AB & DB 11701; AULS 2595 | 6 | NE |
| <i>Clausena heptaphylla</i> (Roxb. ex DC.) Wight & Arn. ex Steud. | Koronful (B) | Apr.–July | AB & DB 11634; AULS 2576 | 11 | NE |
| <i>Micromelum integerrimum</i> (Roxb. ex DC.) Wight & Arn. ex M.Roem. | Not known | Feb.–Sept. | AB & DB 11527; AULS 2656 | 16 | NE |
| Sabiaceae Blume | | | | | |
| <i>Meliosma pinnata</i> (Roxb.) Maxim. | Not known | May–Oct. | AB & DB 11598; AULS 2679 | 1 | NE |
| Salicaceae Mirb. | | | | | |
| <i>Gynocardia odorata</i> Roxb. | Bonsha, Bandarpele (A), Chhalmogra (H) | Jan.–Aug. | AB & DB 11494; AULS 2649 | near Bhaluk Nala at 10 | NE |
| Sapindaceae Juss. | | | | | |
| <i>Sapindus mukorossi</i> Gaertn. | Manishal (A), Ritha (B), Sukathaiphang (Di) | May–Nov. | AB & DB 11597; AULS 2658 | 1 | NE |
| Sapotaceae Juss. | | | | | |
| <i>Mimusops elengi</i> L. | Bakul (A) | May–Sept. | AB & DB 11596; AULS 2632 | 1 | NE |
| Theaceae Mirb. ex Ker Gawl. | | | | | |
| <i>Schima wallichii</i> (DC.) Korth. | Makria, Noga-bhe (A), Chilauni (B) | Apr.–Dec. | AB & DB 11422; AULS 2682 | 5 | NE |
| Ulmaceae Mirbel | | | | | |
| <i>Trema orientalis</i> Blume | Jiban (B), Gunpowder tree (E) | Apr.–Dec. | AB & DB 11528; AULS 2579 | 16 | NE |
| Urticaceae Juss. | | | | | |
| <i>Oreocnide integrifolia</i> (Gaudich.) Miq. | Ban rhea, Bon rhea (A), Wild Rhea (E), | March–Sept. | AB & DB 11387, 11609; AULS 2589 | 3, 11 | NE |
| Verbenaceae J.St.-Hil. | | | | | |
| <i>Gmelina arborea</i> Roxb. ex Sm. | Gamari (A), Gamhar (H) | March–June | AB & DB 11589; AULS 2618 | 11, 6 | NE |
| <i>Tectona grandis</i> L.f. | Segun (A) | Aug.–Jan. | AB 11588; AULS 2626 | 12 | NE |
| Monocotyledons | | | | | |
| Arecaceae C.H. Schultz | | | | | |
| <i>Areca catechu</i> L. | Tamul (A), Supari (B), Betel nut (E) | Apr.–Sept. | AB & DB 11561; AULS 2636 | 15 | NE |
| <i>Caryota urens</i> L. | Chewa gach (A), Fishtail Palm (E) | June-Sept. | AB & DB 11738; AULS 2652 | 1 | LC |
| <i>Licuala peltata</i> Roxb. ex Buch.-Ham. | Chatta-pat (A) | June–Sept. | AB & DB 11498; AULS 2581 | near Bhaluk Nala at 10 | NE |
| Poaceae Barnhart | | | | | |
| <i>Bambusa balcooa</i> Roxb. | Bhuluka banh (A), Guadua Bamboo (E) | Jan.–Aug. | AB & DB 11773; AULS 2586 | 6 | NE |
| <i>Bambusa cacharensis</i> R.B. Majumdar | Betua (B) | Apr.–Nov. | AB & DB 11355, 11356, 11664; AULS 2577 | 5 (way to 1), near Bhaluk Nala at 10 | NE |
| <i>Bambusa tulda</i> Roxb. | Bijuli banh (A) | Jan.–June | AB & DB 11774; AULS 2584 | 11 | NE |
| <i>Bambusa vulgaris</i> Schrad. ex J.C.Wendl. | Tansti banh (A) | Not seen | AB & DB 11772; AULS 2639 | near Bhaluk Nala at 10 | NE |

Table 2. *Continued.*

| Scientific name | Vernacular name | Phenology | Voucher specimens | Occurrence | Threat status |
|--|-----------------|-----------|---------------------------------|----------------------------------|---------------|
| <i>Dendrocalamus hamiltonii</i> Nees & Arn. ex Munro | Kako banh (A) | Oct.–May | AB & DB 11354, 11680; AULS 2655 | way to 1, near Bhaluk Nala at 10 | NE |

Table 3. List of lianas with their vernacular name, phenology, collection number, occurrence and threat status. Abbreviations, languages: A = Assamese, B = Bengali, E = English, H = Hindi, Kh = Khasi, Mik. = Mikir; IUCN status: LC = Least Concern, NE = Not Evaluated. Occurrence: see Table 1 for a key to locations.

| Scientific name | Vernacular name | Phenology | Voucher specimens | Occurrence | Threat status |
|---|---|-------------|-----------------------------------|----------------------------|---------------|
| Dicotyledons | | | | | |
| Acanthaceae Juss. | | | | | |
| <i>Thunbergia grandiflora</i> Roxb. | Kukua loti (A), Neel lota (B) | Sept.–Jan. | AB 11421, 11693; AULS 2603 | 2, 5 | NE |
| Anacardiaceae R.Br. | | | | | |
| <i>Pegia nitida</i> Colebr. | Not known | Jan.–July | AB & DB 11420; AULS 2622 | 5 | NE |
| Apocynaceae Juss. | | | | | |
| <i>Anodendron paniculatum</i> A. DC. | Not known | Apr.–Dec. | AB & DB 11612; AULS 2642 | 11 | NE |
| <i>Hoya globulosa</i> Hook.f. | Not known | Apr.–July | AB & DB 11460; AULS 2685 | 10 | NE |
| Combretaceae R.Br. | | | | | |
| <i>Combretum indicum</i> (L.) DeFilipps | Not known | March–Nov. | AB & DB 11742; AULS 2662 | 11 | NE |
| <i>Combretum pilosum</i> Roxb. | Not known | Dec.–March | AB & DB 11579; AULS 2673 | 3 | NE |
| <i>Combretum wallichii</i> DC. var. <i>deciduum</i> Gang. & T. Chakrab. | Not known | March–Nov. | AB & DB 11426; AULS 2606 | 5 | NE |
| Convolvulaceae Juss. | | | | | |
| <i>Argyreia nervosa</i> (Burm.f.) Bojer | Bichtarak, Goguli (B), Jomang-pi-danok-soh-ring-kang (Mik.) | Oct.–March | AB 11687; AULS 2620 | 2 | NE |
| Fabaceae Lindl. | | | | | |
| <i>Entada gigas</i> (L.) Fawc. & Rendle | Not known | March–Oct. | AB & DB 11417; AULS 2650 | 5 | NE |
| <i>Mucuna imbricata</i> (Roxb.) DC. | Makori ghila (A), Horse eyebean (E) | June–Nov. | AB 11503; AULS 2638 | Kalaincherra (through 8) | LC |
| <i>Phanera glauca</i> Wall. ex Benth. | Not known | Apr.–Sept. | AB 11358, 11694, 11512; AULS 2664 | 2 | LC |
| <i>Phanera ornata</i> Kurz var. <i>kerrii</i> (Gagnep.) K.Larsen & S.S.Larsen | Not known | May | AB 11691; AULS 2671 | 2 | NE |
| <i>Phanera scandens</i> (L.) Raf. | Snake Climber, Monkey ladders (E), Nagbeli (H) | Sept.–Dec. | AB & DB 11534; AULS 2609 | 16 | NE |
| <i>Phanera vahlii</i> (Wight & Arn.) Benth. | Kanchan lota, Nak-kati-lewa (A) | Apr.–Jan. | AB 11495, 11682; AULS 2611 | near Bhaluk Nala at 10, 12 | NE |
| <i>Phanera wallichii</i> Thoth. | Not known | Apr.–July | AB & DB 11601; AULS 2614 | 11 | LC |
| <i>Pueraria montana</i> (Lour.) Merr. var. <i>lobata</i> (Willd.) Maesen & S. M.Almeida ex Sanjappa & Predeep | Kudzu (H) | July–Oct. | AB 11696; AULS 2633 | 2 | NE |
| Hernandiaceae Blume | | | | | |
| <i>Illigera khasiana</i> C.B. Clarke | Kerkerilata (A) | August–Apr. | AB & DB 11427; AULS 2641 | 5 | NE |
| Malvaceae Juss. | | | | | |
| <i>Byttneria aspera</i> Colebr. ex Wall. | Not known | Mar.–Oct. | AB & DB 11463; AULS 2634 | 10 | NE |
| Menispermaceae Juss. | | | | | |
| <i>Tinospora crispa</i> Miers. | Gulanha (B), Heartleaf moonseed (E) | Mar.–Oct. | AB & DB 11523; AULS 2666 | 16 | NE |
| Moraceae Gaudich. | | | | | |
| <i>Poikilospermum suaveolens</i> (Blume) Merr. | Jog-kag, kamlata (B) | Apr.–June | AB & DB 11478; AULS 2604 | 10 | NE |
| Piperaceae Giseke. | | | | | |
| <i>Piper nigrum</i> L. | Jaluk (A), Black pepper (E), Kali mirch (H) | June–Oct. | AB & DB 11713; AULS 2669 | 1 | NE |
| Sabiaceae Blume | | | | | |
| <i>Sabia lanceolata</i> Colebr. | Not known | Oct.–Dec. | AB & DB 11369, 11430; AULS 2691 | 4, 5 | NE |
| Monocotyledons | | | | | |
| Smilacaceae Ventenat | | | | | |
| <i>Smilax zeylanica</i> L. | Kumarika (B) | June–Dec. | AB 11535; AULS 2692 | 12 | NE |

Combretaceae (3 spp.) and Apocynaceae (2 spp.) (Fig. 5). The rest of the 9 families had a single species recorded. The dominant genera were *Phenera* (5 spp.) and *Combretum* (3 spp.). Among the monocotyledonous lianas, a single species (of Smilacaceae) was recorded. Photographs of several species are shown in Figs 8–11.

Common timber yielding trees recorded in the sanctuary were *Artocarpus chama* Buch.-Ham., *A. heterophyllus* Lam., *Azadirachta indica* Juss., *Gmelina arborea* Roxb. ex Sm., *Melia azedarach* L., *Mimusops elengi* L., *Shorea robusta* C.F.Gaertn., *Tectona grandis* L.f., *Terminalia chebula* Retz., and *Toona ciliata* M. Roem. Common in the BWS were edible species of fruit trees: *Aegle marmelos* (L.) Correa, *Carica papaya* L., *Dillenia indica* L., *Elaeocarpus floribundus* Blume, *Mangifera indica* L., *Phyllanthus emblica* L., *Syzygium cumini* (L.) Skeels, *Tamarindus indica* L., and *Zizyphus jujuba* Mill. Other common species of the sanctuary were *Bombax ceiba* L., *Cinnamomum tamala* (Buch.-Ham.) T.Nees & C.H.Eberm., *Ficus benghalensis* L., *F. religiosa* L., *Murraya koenigii* (L.) Spreng., and *Oroxylum indicum* (L.) Kurz. Among the bamboos, *Bambusa vulgaris* Schrad. ex J.C.Wendl. and *Dendrocalamus hamiltonii* Nees & Arn. ex Munro were common in the forest areas. Dominant palm species of the sanctuary were *Areca catechu* L., *Caryota urens* L., and *Licuala peltata* Roxb. ex Buch.-Ham.

Among the lianas, some commonly occurring species recorded were *Combretum indicum* (L.) DeFilipps, *Entada gigas* (L.) Fawc. & Rendle, *Phanera glauca* Wall. ex Benth., *Thunbergia grandiflora* Roxb., and *Tinospora crispa* Miers.

Flowering and fruiting phenology (Figs 6, 7). For trees, the flowering was mostly in March (pre-monsoon) to August (post-monsoon), but peaked in June; the peak in fruit production was in the late dry season (August to December). For lianas, flowering peaked in April to July (pre-monsoon and monsoon) and fruit production peaked in the late dry season (August to December).

IUCN Red List categories. Of all species recorded, only 1 tree species (Vulnerable) is threatened; all other species are categorized as Least Concern, Data Deficient, or are Not Evaluated.

Discussion

Our results indicate a rich flora in the BWS, possibly due to the region's physiography of undulating hills and hillocks and tropical to subtropical climate.

Recently, Barbhuiya (2013a) recorded several species from the BWS that we failed to find. These are: *Actinodaphne obovata* (Nees) Blume, *Balakata bacata* (Roxb.) Esser, *Catunaregam spinosa* (Thunb.) Tirveng., *Cinnamomum bejolghota* (Buch.-Ham.) Sweet, *Croton joufra* Roxb., *Meliosma simplicifolia* (Roxb.) Walp., *Prismatomeris tetrandra* K. Schum., and *Toxicodendron bimannii*. Some earlier collections were also found in the herbarium of Botanical Survey of




Figure 5. Dominant liana families with number of species and genera.

India at ASSAM, which could also not be traced in the field. These are: *Abroma augustum* (L.) L.f., *Archidendron clypearia* (Jack) I.C. Nielsen, *Camellia oleifera* C. Abel, *Dalbergia stipulacea* Roxb., *Dendrocnide sinuata* (Blume) Chew, *Hodgsonia macrocarpa* (Blume) Cong., *Ocotea lancifolia* Mez, *Sterculia villosa* Roxb. ex Sm., and *Styrax serrulatus* Roxb. This might be due to their localized distribution or their rarity in the sanctuary. Additional fieldwork is needed to locate these species in BWS.

Although, most parts of the BWS is still virgin, human habitations in and around the sanctuary exist (8 in East Block; 19 in West Block; Anonymous 2006). With an increasing population, villagers' demands on the forest and its products are increasing with the collection of bamboos, canes, and timber. Clearing of forests for the cultivation of areca nut, beetle leaf, pineapple, and banana is leading to a loss of biodiversity at the forest margins. Massive habitat loss in the buffer zone also




Figure 6. Phenology of the recorded tree species.




Figure 7. Phenology of the recorded liana species.




Figure 8. A. *Byttneria aspera*. B. *Piper nigrum*. C. *Phanera vahlii*. D. *Anodendron paniculatum*. E. *Phanera glauca*.




Figure 9. A. *Hoya globulosa*. B. *Entada gigas*. C. *Pueraria montana* var. *lobata* (Front fruiting twig). D. *Poikilospermum suaveolens*. E. *Smilax zeylanica*. F. *Thunbergia grandiflora*.




Figure 10. **A.** *Ardisia paniculata*. **B.** *Callicarpa arborea*. **C.** *Glochidion lanceolarium*. **D.** *Ehretia acuminata*. **E.** *Gynocardia odorata*. **F.** *Duabanga grandiflora*.




Figure 11. A. *Oreocnide integrifolia*. B. *Saurauia roxburghii*. C. *Saurauia cerea*. D. *Spondias pinnata*. E. *Sterculia versicolor*. F. *Viburnum cylindricum*.

results in species destructions. Other human activities, like road construction, cutting of forests for *jhum* (shifting) cultivation, areca nut plantation etc., along with some natural phenomena like landslides due to heavy rain, thunderstorms, and hailstorms, are also serious threats to biodiversity. Therefore, there is an urgent need for developing and implementing conservation strategies for the BWS.

Acknowledgements

We extend our sincere thanks to the Divisional Forest Officer, Cachar Division, Assam for granting permission to carry out this study. Special thanks are due to Mr D.J. Nath, Forest Range Officer of Kalain Forest Division, for his kind help. We are also thankful to the Head, Botanical Survey of India (ASSAM, and CAL) for permitting us to consult the Institutes' herbaria and libraries. The GUBH herbarium is also gratefully acknowledged. Our gratitude goes to the Dean, School of Life Sciences, Assam University, Silchar and the Head, Department of Life Science and Bioinformatics, Assam University, Silchar for providing facilities. The first author (AB) acknowledges the University Grant Commission (UGC), New Delhi for financial support under the UGC-MRP scheme.

Authors' Contributions

AB carried out fieldwork for collecting specimens, identified the specimens, and drafted the manuscript. DB also took part in field and coordinated the work. Both authors interpreted the results, as well as read and approved the final content of the manuscript.

References

- Anonymous (2006) Management Plan of Barail Wildlife Sanctuary. Cachar and Karimganj Forest Division, Department of Environment and Forests, Government of Assam, Silchar, 112 pp.
- Agarwal NK, Borah A (2001) On the biodiversity of Bhairab Hills of Bongaigaon district of Assam. Part I. Flora. *Journal of Economic and Taxonomic Botany* 25 (2): 247–252.
- APG III (2009) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* 161: 105–121. <https://doi.org/10.1111/j.10958339.2009.00996.x>
- Barbhuiya AR, Sharma GD, Arunachalam A, Deb S (2009) Diversity and conservation of medicinal plants in Barak Valley, northeast India. *Indian Journal Traditional Knowledge* 8 (2): 169–175.
- Barbhuiya HA (2013) *Toxicodendron bimannii* (Anacardiaceae), a new species from Assam, India. *NeBio* 4 (3): 1–5.
- Barbhuiya HA (2013a) Study and Assessment of Threatened and Endemic Vascular Plants of Southern Assam. Ph.D. thesis, Assam University, Silchar, 338 pp.
- Barbhuiya HA (2014) Forty new additions to the angiospermic flora of Assam, India. *India Journal on New Biological Reports* 3 (3): 277–287.
- Barbhuiya HA, Deori C, Dutta BK, Das AKr, Baishya AK, Schuiteman A (2012) *Ornithochilus cacharensis* (Orchidaceae), a new species from India (Assam). *Kew Bulletin* 67: 511–516. <https://doi.org/10.1007/s12225-012-9381-5>
- Barbhuiya HA, Dutta BK, Das AKr, Baishya AK (2012a) Rediscovery of *Ardisia keenanii* C.B. Clarke (Myrsinaceae) from Assam, India. *Taiwania* 57 (1): 82–84. [https://doi.org/10.6165/tai.2012.57\(1\).82](https://doi.org/10.6165/tai.2012.57(1).82)
- Barbhuiya HA, Dutta BK, Das AKr, Baishya AK (2013) An annotated checklist of the grasses (Poaceae) of southern Assam. *Check List* 9 (5): 980–986. <https://doi.org/10.15560/9.5.980>
- Barbhuiya HA, Dutta BK, Das AK and Baishya AK (2014) The family Rubiaceae in southern Assam with special reference to endemic and rediscovered plant taxa. *Journal of Threatened Taxa* 6(4): 5649–5659. doi: <http://dx.doi.org/10.11609/JoTT.03117.5649-59>
- Barooah C, Ahmed I (2014) Plant diversity of Assam—a checklist of angiosperms and gymnosperms. Assam Science Technology and Environment Council, Guwahati, 599 pp.
- Barooah C, Mahanta PK (2006) Aquatic angiosperms of Biswanath Chariali, Assam. *Indian Journal of Forestry* 29 (3): 307–318.
- Barua I (1992) Systematic study of the angiosperms of Kamrup district, Assam. Ph.D. thesis, Gauhati University, Guwahati, 126 pp.
- Baruah MK, Dutta Choudhury M (2015) Flora of Barak Valley, Assam with their Economic Utility, Vol. 2. Herbaceous Flora of Cachar District. Regency Publications, New Delhi, 382 pp.
- Baruah MK, Dutta Choudhury M, Sarma GC (2006) Ethno-medicinal plants used by the Khasi tribe of Cachar district, Assam. *Journal of Economic and Taxonomic Botany* 30 (Suppl.): 110–114.
- Bellefontaine R, Petit S, Pain-Orcet M, Deleporte P, Bertault JG (2002) Trees Outside Forests: Towards Better Awareness. Food and Agriculture Organization, Rome, Conservation Guide 35: 218 pp.
- Bongers F, Schnitzer SA, Traore D (2002) The importance of lianas and consequences for forest management in West Africa. *Bioterre: Revue Internationale des Sciences de la Vie et de la Terre (Special Issue)*: 59–70.
- Brandis D (1906) Indian Trees. Bishen Singh Mahendra Pal Singh, Dehradun, 767 pp.
- Champion HG, Seth SK (1968) A Revised Survey of the Forest Types of India. Government of India Press, Delhi, 404 pp.
- Chaudhuri AB, Naithani HB (1985) Lianas, climber and shrubby climbers. Part III. International Book Distributors, Dehradun, 89 pp.
- Choudhury S (2005) Assam's Flora. Present Status of Vascular Plants. Assam Science, Technology and Environment Council, Guwahati, 368 pp.
- Choudhury S, Katak SK, Barua IC (1994) Floristic analysis of angiosperms of Kamrup district, Assam. *Journal of Economic and Taxonomic Botany* 18 (3): 697–703.
- Darlong L, Bhattacharyya D (2011) *Globba orixensis* Roxb. (Zingiberaceae)—a new record for the flora of Assam, India. *Journal of Bombay Natural History Society* 108 (2): 153–154.
- Darlong L, Das S, Bhattacharyya D (2013) *Drosera burmanni* Vahl (Droseraceae): a common species suffering rapid declines in Barak Valley, Assam. *NeBio* 4 (5): 7–9.
- Das AK, Dutta BK, Sharma GD (2006) Study of medicinal plants used by the different tribes/communities of Cachar district, Assam—I. *Journal of Economic Taxonomic Botany* 30 (Suppl.): 294–309.
- Das AK, Dutta BK, Sharma GD (2007) Study of the tea garden flora of Barak Valley, Assam, India. *Journal of Economic Taxonomic Botany* 31 (4): 858–866.
- Das AK, Dutta BK, Sharma GD (2008) Study of non-conventional edible plants used by different communities of Barak Valley, Assam (N.E. India). *Journal of Economic Taxonomic Botany* 32 (Suppl.): 327–333.
- Das PS, Dutta Choudhury M (2003) A survey on non-conventional food plants of southern Assam. *Journal of Economic and Taxonomic Botany* 27 (2): 416–420.
- Das PS, Dutta Choudhury M, Dutta BK (2013) Flora of Barak Valley, Assam with their Economic Utility, Vol. 1. Herbaceous Flora. Regency Publications, New Delhi, 496 pp.
- Das S (1942) Floristic of Assam—a preliminary sketch. 150th Anniversary Volume. Royal Botanic Garden, Calcutta, 156 pp.
- Devi M, Bhattacharyya D (2013) Flowering occurrences of *Bambusa cacharensis* in Barak Valley, Assam, India since its type collection: a note. *Bamboo Science and Culture: the Journal of the American*

- Bamboo Society 23 (1): 41–44.
- Devi M, Bhattacharyya D (2013a) Gregarious flowering and mast seeding of a climbing bamboo *Melocalamus compactiflorus* in Barak Valley, Assam. *NeBIO* 4 (5): 43–45.
- Devi M, Bhattacharyya D (2013b) *Panicum laxum* (Poaceae): An addition to flora of India. *Rheedea* 23 (2): 128–131.
- Devi M, Bhattacharyya D (2014) Recent flowering of six species of bamboos (Poaceae: Bambusoideae) in northeastern India with identification keys. *Rheedea* 24 (1): 29–45.
- Devi M, Bhattacharyya D (2016) *Themeda quadrivalvis* (L.) Kuntze (Poaceae): A New Record for Assam, India. *Indian Forester* 142 (3): 302–303.
- Devi M, Darlong L, Bhattacharyya D (2014) *Dendrocalamus longispatus* (Poaceae: Bambusoideae) in sporadic bloom in Tripura, India. *NeBIO* 5 (1): 88–89.
- Dutt AK, Dutta TK, Gupta KK (1974) Tentative accounting of the forest flora of North Cachar Hills and Barail range: enumeration of species – dicotyledons, Polypetalae. *Indian Forester* 100 (1) 60–76.
- Grubb PJ (1977) Control of forest growth and distribution on wet tropical mountains with special reference to mineral nutrition. *Annual Review of Ecology and Systematic* 8: 83–107.
- IPNI (2012) The International Plant Names Index. <http://www.ipni.org>. Accessed on: 2016-11-12.
- Islam M (1986) A contribution to the flora of the greatest river island Majuli, Jorhat, Assam-II. *Journal of Economic and Taxonomic Botany* 8 (1): 77–105.
- Islam M (1990) The Flora of Majuli. Bishen Singh Mahendra Pal Singh, Dehradun, 436 pp.
- Islam M (1990a) A contribution to the flora of the greatest river island Majuli, Jorhat, Assam—III. *Journal of Economic and Taxonomic Botany* 14 (1): 27–31.
- Islam M (1991) Exotic flora of Assam. *Journal of Economic and Taxonomic Botany* 15 (2): 391–407.
- IUCN (2017) The IUCN Red List of Threatened Species. Version 2017-1. <http://www.iucnredlist.org>. Accessed on: 2017-4-2.
- Jain SK, Rao RR (1977) *Handbook of Field and Herbarium Methods*. Today and Tomorrow's Printers and Publishers, New Delhi, 157 pp.
- Kanjilal UN, Kanjilal PC, Das A, Dey RN, Bor NL (1934–1940) *Flora of Assam*. Vols. 1–5. Government of Assam, Shillong.
- Khandekar V, Srivastava A (2014) Ecosystem biodiversity of India. In: Grillo O (Ed.) *Biodiversity— the Dynamic Balance of the Planet*. InTech, Rijeka, 378 pp. <https://doi.org/10.5772/57015>
- Kress WJ, Mood JD, Sabu M, Prince LM, Dey S, Sanoj E (2010) *Larsenianthus*, a new Asian genus of gingers (Zingiberaceae) with four species. *PhytoKeys* 1: 15–32. <https://doi.org/10.3897/phytokeys.1.658>
- Nath A, Maiti GG (2003) Ethnobotany of Barak Valley (Southern Assam) with special reference to folk medicine. *Journal of Economic and Taxonomic Botany* 27 (4): 964–971.
- Rao AS, Rabha LC (1966) Contribution to the botany of Kamrup district (southern part), Assam. *Bulletin of the Botanical Survey of India* 8: 296–303. https://doi.org/10.20324/nelumbo/v8/1966/76211_
- Rao AS, Verma DM (1970) Materials towards a monocot flora of Assam (Hydrocharitaceae and Burmanniaceae). *Bulletin of the Botanical Survey of India* 12 (1–4): 139–143. <https://doi.org/10.20324/nelumbo/v12/1970/75850>
- Rao AS, Verma DM (1972) Materials towards a monocot flora of Assam II (Zingiberaceae and Marantaceae). *Bulletin of the Botanical Survey of India* 14 (1–4): 114–143. <https://doi.org/10.20324/nelumbo/v14/1972/75253>
- Rao AS, Verma DM (1973) Materials towards a monocot flora of Assam III (Taccaceae, Dioscoreaceae and Stemonaceae). *Bulletin of the Botanical Survey of India* 15 (3–4): 189–209. https://doi.org/10.20324/nelumbo/v15/1973/75162_
- Rao AS, Verma DM (1976) Materials towards a monocot flora Assam IV (Pontederiaceae, Xyridaceae & Commelinaceae). *Bulletin of the Botanical Survey of India* 16 (1–4): 1–20. https://doi.org/10.20324/nelumbo/v16/1974/75003_
- Rout J, Sajem AL, Nath M (2012) Medicinal plants of North Cachar Hills district of Assam used by the Dimasa tribe. *National Journal of Traditional Knowledge* 11 (3): 520–527.
- Sarma SK (1990) Floristic composition and phytosociological studies of a herbaceous community at Dibrugarh. *Indian Journal of Forestry* 13 (4): 273–280.
- Sarma SK, Devi B, Bhattacharyya DK (2006) Ethnomedicinal uses of plants by the Sarania tribe of Nalbari district of Assam. *Journal of Economic and Taxonomic Botany* 30 (Suppl.): 133–139.
- Sharma GD, Bhattacharjee S, Sinha M, Das AK (2002) Status of plant bio-diversity of Cachar district and its conservation. *Journal of Economic and Taxonomic Botany* 26: 94–101.
- Singh HB, Subramaniam B (2008) *Field Manual on Herbarium Techniques*. National Institute of Science Communication and Information Resources, Council of Scientific & Industrial Research (CSIR), New Delhi, 298 pp.
- The Plant List (2013) The Plant List. Version 1.1. <http://www.theplantlist.org/>. Accessed on: 2016-3-20.
- Thiers B (2011) *Index Herbariorum: A Global Directory of Public Herbaria and Associated Staff*. New York Botanical Garden's Virtual Herbarium. <http://sciweb.nybg.org/science2/IndexHerbariorum.asp>.
- Tropicos (2017) Tropicos. <http://www.tropicos.org/Home.aspx>. Accessed on: 2017-4-5.